

Chemical Composition, Antioxidant Capacity, Acetyl- and Butyrylcholinesterase Inhibitory Activities of the Essential Oil of *Thymus haussknechtii* Velen.

Handan G. Sevindik¹, Temel Özek², K. Özden Yerdelen³, Mehmet Önal⁴, Hilal Özbek^{1*}, Zühal Güvenalp¹ and L. Ömür Demirezer⁵

¹Department of Pharmacognosy, Faculty of Pharmacy, Atatürk University, Erzurum, 25240, Türkiye

²Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, Eskişehir, 26470, Türkiye

³Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Atatürk University, Erzurum, 25240, Türkiye

⁴Non-wood Products Chief Engineering, Regional Directorate of Forestry, Erzurum, 25050, Türkiye

⁵Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, Ankara, 06100, Türkiye

(Received January 24, 2015; Revised August 11, 2015; Accepted August 15, 2015)

Abstract: The chemical composition of the essential oil from the aerial parts of *Thymus haussknechtii* Velen. was analyzed by using gas chromatography (GC-FID) and gas chromatography-mass spectrometry (GC-MS). The major component of the essential oil was thymol (52.2%). Total phenolic content of the essential oil was determined as 132.9 µg gallic acid equivalent. The antioxidant capacity was evaluated by DPPH free radical, superoxide anion radical and hydrogen peroxide scavenging activities along with ferrous ion-chelating power test, ABTS radical cation decolorization assay and ferric thiocyanate methods. In addition to antioxidant activity, anticholinesterase activity of the essential oil was also evaluated. It exhibited inhibitory activities on AChE and BuChE which play an important role in Alzheimer's disease, along with significant antioxidant activity.

Keywords: *Thymus haussknechtii*; essential oil; antioxidant activity; acetylcholinesterase; butyrylcholinesterase. © 2016 ACG Publications. All rights reserved.

1. Plant Source

The genus *Thymus* (Lamiaceae) is represented by 43 species in Turkey [1-4] and the biological activity studies showed that the essential oil of some *Thymus* species indicated antioxidant, antimicrobial, insecticidal [5-8], antiprotective [9], antibacterial [10-11] activities.

The aerial parts of *Thymus haussknechtii* Velen. were collected from Mount Kop in September 2013 (2030 m) and identified by Mehmet Önal from Regional Directorate of Forestry, Türkiye. The

* Corresponding author: E- Mail: ozbek@atauni.edu.tr (H. Özbek), Phone +90-442-2315227.

voucher specimen (AUEF 1001) has been deposited in the herbarium of the Faculty of Pharmacy, Atatürk University, Erzurum, Türkiye.

2. Previous Studies

A few study have been carried out on the chemical composition of the essential oil of *T. haussknechtii* which were collected from different places of Turkey and exhibited different chemical profiling [7, 12-14]. The main constituents of the oil of *T. haussknechtii* collected from Erzincan were reported as 1,8-cineole (23.6%), *trans*-verbenol (6.6%), camphor (6.12%) and caryophyllene oxide (6%). Furthermore, its antioxidant activity was determined by means of the DPPH radical-scavenging method and at 1000 µg/mL concentration of the essential oil 35.11 ± 0.22 % DPPH was scavenged [7]. The essential oil of *T. haussknechtii* collected from Elazığ province was found to possess linalool (19.9%) and borneol (10.3%) as major components [13]. Finally, 1,8-cineole (23.6%) was designated as the main constituent in the oil of *T. haussknechtii* collected from Elazığ Harput-Ankuzubaba Mountain [14].

3. Present Study

Chemical composition of the essential oil: Hydrodistillation of the aerial parts of *T. haussknechtii* collected from Kop Mount, gave light yellow oil with 1.34% (w/w) yield. The essential oil composition was analyzed by GC-FID and GC-MS. Forty nine components constituting 99.6% of the essential oil were identified. Thymol was the main constituent of *T. haussknechtii* essential oil with 52.2% of the total oil composition. All components of the essential oil identified by GC and GC-MS analyses were listed in Table 1 along with their relative retention indices and their percentage composition.

GC analysis of the essential oil was carried out on Agilent 5975 GC-MSD system and Agilent 6890N GC system with flame ionization detector (FID) and a HP-Innowax FSC column (60 m x 0.25 mm, 0.25 µm film thickness). Helium was used as a carrier gas at 0.8 mL/min. The injector and detector temperatures were 250°C and 300°C, respectively. GC oven temperature was at 60°C for 10 min, then programmed to 220°C at a rate of 4°C/min and kept in this temperature for 10 min and finally programmed to 240°C at a rate of 1°C/min with a final hold time of 80 min. The split ratio was 40:1. Mass spectrums were taken at 70 eV and mass range was from *m/z* 35 to 450. In order to obtain same elution order with GC-MS, simultaneous injection was done by using the same column and appropriate operational conditions.

In consideration of previous studies, the major constituents of *T. haussknechtii* oils from different localities were found to be 1,8-cineole, *trans*-verbenol, camphor, caryophyllene oxide, linalool and borneol in the range of 6-23.6%. However, the main component of our essential oil was thymol with a high amount (52.2%). These results indicates that geographical location, collection season and environmental factors may be the reason of these variations in the essential oil composition with important quantitative differences [6-7,12-14].

Total phenolic content of the essential oil: Total phenolic content of the essential oil was determined according to the Folin-Ciocalteu's method [15] using gallic acid as a standard phenolic compound. Total phenolic content of the essential oil was calculated by using the equation that was obtained from the standart gallic acid graph (Figure S1). 132.9 ± 0.2 µg gallic acid equivalent of phenolic compounds were detected in 1 mg of the essential oil.

Antioxidant activity tests: Antioxidant activity was evaluated on six tests: DPPH free radical scavenging activity [16], ferrous ion-chelating power test [17], ABTS radical cation decolorization assay [18], superoxide anion radical scavenging activity [19], total antioxidant activity by ferric thiocyanate method [20], hydrogen peroxide scavenging activity [21], which were carried out as described in the literatures. The scavenging effect of the essential oil of *T. haussknechtii*, thymol as the major component of the essential oil and reference compound trolox on DPPH, ABTS, H₂O₂ and superoxide radical were evaluated.

Table 1. Chemical composition of the essential oil of *T. haussknechtii*.

Compound	RRI^a	%^b
Tricyclene	1014	t ^c
α -Pinene	1032	0.7
α -Thujene	1035	0.7
Camphene	1076	0.5
β -Pinene	1118	0.3
Sabinene	1132	0.1
δ -3-Carene	1159	t
Myrcene	1174	0.8
α -Terpinene	1188	0.9
Limonene	1203	0.3
1,8-Cineole	1213	4.6
γ -Terpinene	1255	6.6
3-Octanone	1266	0.1
<i>p</i> -Cymene	1280	16.6
Terpinolene	1290	0.2
3-Octanol	1393	0.1
α , <i>p</i> -Dimethylstyrene	1452	t
1-Octen-3-ol	1452	0.4
<i>trans</i> -Sabinene hydrate	1474	0.3
α -Copaene	1497	t
β -Bourbonene	1535	0.1
<i>cis</i> -Sabinene hydrate acetate	1542	0.2
<i>trans-p</i> -Menth-2-en-1-ol	1571	0.2
β -Ylangene	1589	0.1
Bornyl acetate	1590	0.3
Terpinen-4-ol	1611	1.6
β -Caryophyllene	1612	0.7
<i>cis-p</i> -Menth-2-en-1-ol	1638	t
δ -Terpineol	1682	0.2
<i>trans</i> -Verbenol	1683	0.2
γ -Muurolene	1704	0.1
α -Terpineol	1706	0.7
Borneol	1719	2.2
Germacrene D	1726	0.2
β -Bisabolene	1741	0.4
<i>cis</i> -Piperitol	1758	t
δ -Cadinene	1773	0.1
γ -Cadinene	1776	t
Cumin aldehyde	1802	t
<i>cis</i> -Calamenene	1853	t
<i>p</i> -Cymen-8-ol	1864	0.2
Thymol acetate	1867	0.4
(<i>E</i>)-Nerolidol	2050	0.3
Humulene epoxide-II	2071	1.9
Cumin alcohol	2113	0.1
Isothymol (=2-Isopropyl-4-methyl phenol)	2181	0.2
Thymol	2198	52.2
Isocarvacrol (=4-Isopropyl-2-methyl phenol)	2221	0.2
Carvacrol	2239	3.6
Total	99.6	

^a Relative retention indices (RRI) calculated against *n*-alkanes.

^b The contents (%) of the individual components were calculated based on the peak area (FID response).

^c t: Trace (< 0.1 %).

As shown in Table 2, the essential oil had significant scavenging effect on ABTS, H₂O₂ and superoxide radical at 60 μ g/mL with percentage inhibition of 100, 100 and 77.20%, when compared with that of thymol 97.08, 93.27, 61.72% and trolox 97.89, 81.72, 61.36%, respectively. Fe²⁺ chelating ability of the essential oil and thymol at 60 μ g/mL were determined as 17.33 and 21.42%. The total antioxidant activity of the essential oil and thymol were measured by the inhibition of lipid peroxidation with percentage inhibition of 41.44 and 42.24% which were comparable to that of the

standard compound (44.71%). According to the present study, the essential oil of *T. haussknechtii* was found to be an effective antioxidant according to different *in vitro* assays when it was compared to standard antioxidant compound trolox. The effect was considered to be due to its major component thymol which could play an important role in the antioxidant activity because of its phenolic structure [11].

Table 2. Antioxidant activity of the essential oil of *T. haussknechtii* at 60 µg/mL.

	Inhibition (%) ^a		
	Essential Oil	Thymol	Trolox
ABTS radical cation decolorization activity	100.00 ± 0	97.08 ± 1.01	97.89 ± 0.21
DPPH free radical scavenging activity	3.24 ± 1.81	23.66 ± 0.29	45.92 ± 0.32
Ferrous ion chelating activity	17.33 ± 2.43	21.42 ± 0.68	59.83 ± 0.19
Superoxide anion radical scavenging activity	77.20 ± 1.17	61.72 ± 1.42	61.36 ± 1.03
Total antioxidant activity	41.44 ± 0.52	42.24 ± 1.15	44.71 ± 1.27
H ₂ O ₂ scavenging activity	100.00 ± 0	93.27 ± 0.84	81.72 ± 0.41

^a % inhibition (means ± SD)

The microplate assay for anticholinesterase activity: Inhibitory activities of AChE and BuChE of the essential oil were evaluated by colorimetric Ellman's method [22] with some modifications using commercially available neostigmine bromide as the reference compound [23]. In this study, the essential oil of *T. haussknechtii* showed anticholinesterase activity against acetylcholinesterase with 57.33% and butyrylcholinesterase with 40.11% inhibition at 25 µg/mL concentration (Table 3). The essential oil obtained from *T. haussknechtii* was rich in thymol (52.2%) and thymol exhibited strong acetyl- (83.0%) and butyrylcholinesterase (98.0%) inhibitory activities. Therefore, the anticholinesterase activity of the essential oil could be attributed to its major constituent thymol [24]. We present the first report on anticholinesterase activity of the essential oil of *T. haussknechtii* in this study.

Table 3. *In vitro* AChE and BuChE inhibition of the essential oil of *T. haussknechtii* at 25 µg/mL.

	AChE ^a (%)	BuChE ^b (%)
Essential oil	57.33 ± 3.05	40.11 ± 2.08
Thymol	83.0 ± 0.11	98.0 ± 1.38
Neostigmine bromide	100 ± 0	100 ± 0

^a 50% inhibitory concentration (means ± SD of three experiments) of AChE.

^b 50% inhibitory concentration (means ± SD of three experiments) of BuChE.

It can be concluded that the essential oil of *T. haussknechtii* possess strong antioxidant effect and anticholinesterase activity which may be due to its high phenolic content. These results indicate that *T. haussknechtii* could be a good source for natural antioxidants which were very important in prevention of many disease and protection of health.

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/RNP>

References

- [1] J. Jalas (1982). *Thymus* L., In: Flora of Turkey and East Aegean Islands, ed: P. H. Davis, University Press, Edinburgh, pp.349-382.
- [2] J. Jalas (1988). *Thymus* L., In: Flora of Turkey and East Aegean Islands, eds: P. H. Davis, R. R. Mill and K. Tan, University Press, Edinburgh, pp.209-210.
- [3] H. Duman (2000). *Thymus* L., In: Flora of Turkey and East Aegean Islands, eds: A. Güner, N. Özhatay, T. Ekim and K. H. C. Başer, University Press, Edinburgh, pp.209.
- [4] A. Güner, S. Aslan, T. Ekim, M. Vural and M. T. Babaç (2012). Türkiye Bitkileri Listesi (Damarlı Bitkiler). Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul.

- [5] F. Z. Küçükbay, E. Kuyumcu, S. Çelen, A. D. Azaz, T. Arabaci and B. Yildiz (2013). Chemical Composition and Antimicrobial and Antioxidant Activities of three Turkish Thyme Essential Oils, *J.E.O.B.* **16**, 661-671.
- [6] F. Z. Küçükbay, E. Kuyumcu, S. Çelen, A. D. Azaz and T. Arabacı (2014). Chemical composition of the essential oils of three *Thymus* taxa from Turkey with antimicrobial and antioxidant activities, *Rec. Nat. Prod.* **8**, 110-120.
- [7] S. Celen, A. D. Azaz and F. Z. Kucukbay (2012). Chemical composition, antimicrobial and antioxidant properties of *Thymus haussknechtii* Velen. essential oil, *Journal of Applied Biological Sciences* **6**, 13-18.
- [8] R. Belağziz, R. Harrak, A. Romane, K. Oufdou and M. A. El Alaoui ElFels (2010). Antimicrobial and insecticidal activities of the endemic *Thymus broussonetti* Boiss. and *Thymus maroccanus* Ball., *Rec. Nat. Prod.* **4**, 230-237.
- [9] B. Cetin, S. Cakmakçı and M. Gurses (2013). Anti-Probiotic Effects of Essential Oils from Some Turkish Endemic Thyme Species, *Asian J. Chem.* **25**, 8625-8628.
- [10] R. Kotan, A. Cakir, F. Dadasoglu, T. Aydın, R. Cakmakci, H. Ozer, S. Kordali, E. Mete and N. Dikbas (2010). Antibacterial activities of essential oils and extracts of Turkish *Achillea*, *Satureja* and *Thymus* species against plant pathogenic bacteria, *J. Sci. Food Agric.* **90**, 145-160.
- [11] H. Nazemiyeh, F. Lotfipoor, A. Delazar, S. M. Razavi, S. Asnaashari, N. Kasebi, A. H. Talebpour, L. Nahar and S. D. Sarker (2011). Chemical composition, and antibacterial and free-radical scavenging activities of the essential oils of a citronellol producing new chemotype of *Thymus pubescens* Boiss. & Kotschy ex Celak, *Rec. Nat. Prod.* **5**, 184-192.
- [12] F. Ilisulu Meriçli (1986). Evaluation of the thymol contents of endemic *Thymus* species growing in Turkey, *Doga: Turk Tip ve Eczacilik Dergisi* **10**, 187-200.
- [13] K. H. C. Baser, T. Ozek and G. Tumen (1992). Essential oils of *Thymus cariensis* and *Thymus haussknechtii*, two endemic species in Turkey, *J. Essent. Oil Res.* **4**, 659-661.
- [14] E. Bağcı and K. H. C. Başer (2005). Study of the essential oils of *Thymus haussknechtii* Velen and *Thymus kotschyanus* Boiss. et Hohen var. *kotschyanus* (Lamiaceae) taxa from the eastern Anatolian region in Turkey, *Flavour Fragr. J.* **20**, 199-202.
- [15] K. Slinkard and V. L. Singleton (1977). Total phenol analyses: automation and comparison with manual methods, *Am. J. Enol. Viticul.* **28**, 49-55.
- [16] M. S. Blois (1958). Antioxidant determinations by the use of a stable free radical, *Nature* **26**, 1199-1200.
- [17] T. C. P. Dinis, V. M. C. Madeira and L. M. Almeida (1994). Action of phenolic derivatives (acetoaminophen, salicylate, and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxyl radical scavengers, *Arch. Biochem. Biophys.* **315**, 161-169.
- [18] R. Re, N. Pellegrini, A. Proteggente, A. Pannala, M. Yang and C. Rice-Evans (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay, *Free Radical Bio. Med.* **26**, 1231-1237.
- [19] J. Zhishen, T. Mengcheng and W. Jianming (1999). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals, *Food Chem.* **64**, 555-559.
- [20] H. Mitsuda, K. Yasumoto and K. Iwami (1966). Antioxidative action of indole compounds during the autoxidation of linoleic acid, *Eiyoto Shokuryo* **19**, 210-214.
- [21] R. J. Ruch, S. J. Cheng and J. F. Klaunig (1989). Prevention of cytotoxicity and inhibition of intracellular communication by antioxidant catechins isolated from Chinese green tea, *Carcinogenesis* **10**, 1003-1008.
- [22] G. L. Ellman, D. Courtney, V. Andies and R. M. Featherstone (1961). A new and rapid colorimetric determination of acetylcholinesterase activity, *Biochem. Pharmacol.* **7**, 88-95.
- [23] K. Ö. Yerdelen and E. Tosun (2015). Synthesis, docking and biological evaluation of oxamide and fumaramide analogs as potential AChE and BuChE inhibitors, *Med. Chem. Res.* **24**, 588-602.
- [24] M. Öztürk (2012). Anticholinesterase and antioxidant activities of Savoury (*Satureja thymbra* L.) with identified major terpenes of the essential oil, *Food Chem.* **134**, 48-54.