

Diaquabis(2-bromobenzoato- κ O)bis-(nicotinamide- κ N¹)nickel(II)

Tuncer Hökelek,^{a*} Filiz Yılmaz,^b Barış Tercan,^c F. Elif Özbek^d and Hacali Necefoğlu^d

^aDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey,

^bDepartment of Chemistry, Faculty of Science, Anadolu University, 26470

Yenibağlar, Eskişehir, Turkey, ^cDepartment of Physics, Karabük University, 78050

Karabük, Turkey, and ^dKafkas University, Department of Chemistry, 63100 Kars,

Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 8 June 2009; accepted 8 June 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.034; wR factor = 0.090; data-to-parameter ratio = 16.6.

The title Ni^{II} complex, $[\text{Ni}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2(\text{H}_2\text{O})_2]$, is centrosymmetric. It contains two 2-bromobenzoate (BB) ligands, two nicotinamide (NA) ligands and two water molecules, all of them being monodentate. The four O atoms in the equatorial plane around the Ni atom form a slightly distorted square-planar arrangement, while the slightly distorted octahedral coordination is completed by the two N atoms of the NA ligands in the axial positions. The dihedral angle between the carboxylate group and the adjacent benzene ring is $30.81(17)^\circ$, while the pyridine and benzene rings are oriented at a dihedral angle of $84.66(6)^\circ$. In the crystal structure, $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into a supramolecular structure. A weak $\text{C}-\text{H}\cdots\pi$ interaction is also found.

Related literature

For general background, see: Antolini *et al.* (1982); Bigoli *et al.* (1972); Krishnamachari (1974); Nadzhafov *et al.* (1981); Shnulin *et al.* (1981). For related structures, see: Hökelek *et al.* (2009a,b,c); Özbek *et al.* (2009); Sertçelik *et al.* (2009a,b,c); Tercan *et al.* (2009).



Experimental

Crystal data

$[\text{Ni}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2$

$(\text{H}_2\text{O})_2]$

$M_r = 739.02$

Monoclinic, $P2_1/n$

$a = 7.8851(2)$ Å

$b = 18.2865(4)$ Å

$c = 9.7574(3)$ Å

$\beta = 106.609(2)^\circ$

$V = 1348.23(6)$ Å³

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 3.74$ mm⁻¹

$T = 100$ K

$0.52 \times 0.27 \times 0.22$ mm

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.306$, $T_{\max} = 0.436$

12686 measured reflections

3368 independent reflections

2899 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.090$

$S = 1.06$

3368 reflections

203 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 1.48$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.55$ e Å⁻³

Table 1

Selected bond lengths (Å).

Ni1—O1	2.0806 (16)	Ni1—N1	2.068 (2)
Ni1—O4	2.1012 (17)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H21 \cdots O2 ⁱ	0.82 (4)	2.09 (4)	2.864 (3)	156 (4)
N2—H22 \cdots O3 ⁱⁱ	0.79 (4)	2.22 (4)	2.951 (3)	153 (4)
O4—H41 \cdots O2 ⁱⁱⁱ	0.86 (4)	1.77 (4)	2.612 (2)	165 (5)
O4—H42 \cdots O3 ^{iv}	0.83 (4)	2.09 (4)	2.886 (2)	162 (3)
C9—H9 \cdots Cg1 ^v	0.93	2.89	3.617 (3)	136

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x+2, -y+1, -z$; (iii) $-x+1, -y+1, -z+1$; (iv) $-x+2, -y+1, -z+1$; (v) $-x, -y, -z+1$. Cg1 is the centroid of the C2–C7 ring.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics:

ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

The authors are indebted to Anadolu University and the Medicinal Plants and Medicine Research Centre of Anadolu University, Eskişehir, Turkey, for the use of X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2538).

References

- Antolini, L., Battaglia, L. P., Corradi, A. B., Marcotrigiano, G., Menabue, L., Pellacani, G. C. & Saladini, M. (1982). *Inorg. Chem.* **21**, 1391–1395.
- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst.* **B28**, 962–966.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hökelek, T., Dal, H., Tercan, B., Özbek, F. E. & Necefoğlu, H. (2009a). *Acta Cryst.* **E65**, m466–m467.
- Hökelek, T., Dal, H., Tercan, B., Özbek, F. E. & Necefoğlu, H. (2009b). *Acta Cryst.* **E65**, m481–m482.
- Hökelek, T., Dal, H., Tercan, B., Özbek, F. E. & Necefoğlu, H. (2009c). *Acta Cryst.* **E65**, m533–m534.
- Krishnamachari, K. A. V. R. (1974). *Am. J. Clin. Nutr.* **27**, 108–111.
- Nadzhafov, G. N., Shnulin, A. N. & Mamedov, Kh. S. (1981). *Zh. Strukt. Khim.* **22**, 124–128.
- Özbek, F. E., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009). *Acta Cryst.* **E65**, m341–m342.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009a). *Acta Cryst.* **E65**, m324–m325.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009b). *Acta Cryst.* **E65**, m326–m327.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009c). *Acta Cryst.* **E65**, m389–m390.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shnulin, A. N., Nadzhafov, G. N., Amiraslanov, I. R., Usubaliev, B. T. & Mamedov, Kh. S. (1981). *Koord. Khim.* **7**, 1409–1416.
- Tercan, B., Hökelek, T., Aybirdi, Ö. & Necefoğlu, H. (2009). *Acta Cryst.* **E65**, m109–m110.

supplementary materials

Acta Cryst. (2009). E65, m768-m769 [doi:10.1107/S1600536809021710]

Diaquabis(2-bromobenzoato- κO)bis(nicotinamide- κN^1)nickel(II)

T. Hökelek, F. Yilmaz, B. Tercan, F. E. Özbek and H. Necefoglu

Comment

Transition metal complexes with biochemically active ligands frequently show interesting physical and/or chemical properties, as a result they may find applications in biological systems (Antolini *et al.*, 1982). The structural functions and coordination relationships of the arylcarboxylate ion in transition metal complexes of benzoic acid derivatives change depending on the nature and position of the substituent groups on the benzene ring, the nature of the additional ligand molecule or solvent, and the medium of the synthesis (Nadzhafov *et al.*, 1981; Shnulin *et al.*, 1981). Nicotinamide (NA) is one form of niacin and a deficiency of this vitamin leads to loss of copper from the body, known as pellagra disease. Victims of pellagra show unusually high serum and urinary copper levels (Krishnamachari, 1974). On the other hand, the nicotinic acid derivative *N,N*-diethylnicotinamide (DENA) is an important respiratory stimulant (Bigoli *et al.*, 1972).

The structure determination of the title compound, (I), a nickel complex with two 2-bromobenzoate (BB), two nicotinamide (NA) ligands and two water molecules, was undertaken in order to determine the properties of the ligands and also to compare the results obtained with those reported previously.

Compound (I) is a monomeric complex, with the Ni atom on a centre of symmetry. It contains two BB, two NA ligands and two water molecules (Fig. 1). All ligands are monodentate. The four O atoms (O1, O4, and the symmetry-related atoms, O1', O4') in the equatorial plane around the Ni atom form a slightly distorted square-planar arrangement, while the slightly distorted octahedral coordination is completed by the two N atoms of the NA ligands (N1, N1') in the axial positions (Table 1 and Fig. 1).

The near equality of the C1—O1 [1.260 (3) Å] and C1—O2 [1.258 (3) Å] bonds in the carboxylate group indicates a delocalized bonding arrangement, rather than localized single and double bonds, and may be compared with the corresponding distances: 1.262 (3) and 1.249 (3) Å in [Mn(DENA)₂(C₈H₅O₃)₂(H₂O)₂], (II) (Sertçelik *et al.*, 2009a), 1.263 (4) and 1.249 (4) Å in [Ni(DENA)₂(C₈H₅O₃)₂(H₂O)₂], (III) (Sertçelik *et al.*, 2009b), 1.262 (5) and 1.257 (5) Å in [Co(DENA)₂(C₈H₅O₃)₂(H₂O)₂], (IV) (Sertçelik *et al.*, 2009c), 1.244 (4) and 1.270 (4) Å in [Co(NA)₂(H₂O)₄](C₇H₄FO₂)₂, (V) (Özbek *et al.*, 2009), 1.284 (2), 1.248 (2) and 1.278 (2), 1.241 (2) Å in [Zn(NA)₂(C₈H₈NO₂)₂], (VI) (Tercan *et al.*, 2009), 1.267 (3) and 1.258 (3) Å in [Ni(NA)₂(C₇H₄ClO₂)₂(H₂O)₂], (VII) (Hökelek *et al.*, 2009a), 1.263 (2) and 1.240 (2) Å in [Zn(DENA)₂(C₇H₄BrO₂)₂(H₂O)₂], (VIII) (Hökelek *et al.*, 2009b), and 1.2611 (17) and 1.2396 (19) Å in [Mn(DENA)₂(C₇H₄BrO₂)₂(H₂O)₂], (IX) (Hökelek *et al.*, 2009c). In (I), the average Ni—O bond length is 2.0909 (17) Å and the Ni atom is displaced out of the least-squares plane of the carboxylate group (O1/C1/O2) by -0.595 (1) Å. The dihedral angle between the planar carboxylate group and the benzene ring A (C2—C7) is 30.81 (17)°, while that between rings A and B (N1/C8—C12) is 84.66 (6)°.

In the crystal structure, intermolecular O—H...O and N—H...O hydrogen bonds (Table 2) link the molecules into a supramolecular structure, in which they may be effective in the stabilization of the structure. A weak C—H... π interaction (Table 2) is also found.

Experimental

The title compound was prepared by the reaction of Ni(SO₄).6(H₂O) (1.31 g, 5 mmol) in H₂O (20 ml) and NA (1.22 g, 10 mmol) in H₂O (20 ml) with sodium 2-bromobenzoate (2.23 g, 10 mmol) in H₂O (50 ml). The mixture was filtered and set aside to crystallize at ambient temperature for 2 d, giving blue single crystals.

Refinement

H atoms of water molecule and NH₂ group were located in difference Fourier maps and refined isotropically. The remaining H atoms were positioned geometrically with C—H = 0.93 Å, for aromatic H atoms and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Primed atoms are generated by the symmetry operator (1 - x, 1 - y, 1 - z).

Diaquabis(2-bromobenzoato-κO)bis(nicotinamide-κN¹)nickel(II)

Crystal data

[Ni(C₇H₄BrO₂)₂(C₆H₆N₂O)₂(H₂O)₂]

$M_r = 739.02$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 7.8851$ (2) Å

$b = 18.2865$ (4) Å

$c = 9.7574$ (3) Å

$\beta = 106.609$ (2)°

$V = 1348.23$ (6) Å³

$Z = 2$

$F_{000} = 740$

$D_x = 1.820$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 5380 reflections

$\theta = 2.5$ – 28.4 °

$\mu = 3.74$ mm⁻¹

$T = 100$ K

Block, blue

$0.52 \times 0.27 \times 0.22$ mm

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100$ K

ϕ and ω scans

3368 independent reflections

2899 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\text{max}} = 28.4$ °

$\theta_{\text{min}} = 2.2$ °

Absorption correction: multi-scan
(SADABS; Bruker, 2005) $h = -10 \rightarrow 10$
 $T_{\min} = 0.306$, $T_{\max} = 0.436$ $k = -24 \rightarrow 21$
 12686 measured reflections $l = -12 \rightarrow 13$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.034$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.090$ $w = 1/[\sigma^2(F_o^2) + (0.0538P)^2 + 0.96P]$
 $S = 1.06$ where $P = (F_o^2 + 2F_c^2)/3$
 3368 reflections $(\Delta/\sigma)_{\max} < 0.001$
 203 parameters $\Delta\rho_{\max} = 1.48 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.55 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.40260 (3)	0.212024 (14)	0.17902 (3)	0.02187 (10)
Ni1	0.5000	0.5000	0.5000	0.01048 (11)
O1	0.6201 (2)	0.40537 (9)	0.45455 (18)	0.0137 (3)
O2	0.3946 (2)	0.36790 (10)	0.27275 (19)	0.0162 (4)
O3	0.9542 (2)	0.51033 (10)	0.17931 (19)	0.0184 (4)
O4	0.7554 (2)	0.54252 (10)	0.58888 (19)	0.0149 (4)
H41	0.724 (6)	0.574 (2)	0.643 (4)	0.045 (11)*
H42	0.838 (5)	0.5193 (19)	0.643 (4)	0.024 (8)*
N1	0.5076 (3)	0.54372 (11)	0.3063 (2)	0.0129 (4)
N2	0.8578 (3)	0.57987 (14)	-0.0186 (2)	0.0194 (5)
H21	0.776 (5)	0.6019 (19)	-0.074 (4)	0.034 (10)*
H22	0.935 (5)	0.5647 (18)	-0.048 (4)	0.023 (8)*
C1	0.5546 (3)	0.36682 (13)	0.3452 (2)	0.0126 (4)

supplementary materials

C2	0.6793 (3)	0.31957 (13)	0.2910 (3)	0.0130 (4)
C3	0.6274 (3)	0.25649 (14)	0.2082 (3)	0.0138 (5)
C4	0.7416 (3)	0.22059 (13)	0.1454 (3)	0.0160 (5)
H4	0.7036	0.1793	0.0893	0.019*
C5	0.9119 (3)	0.24655 (15)	0.1666 (3)	0.0176 (5)
H5	0.9876	0.2235	0.1227	0.021*
C6	0.9699 (3)	0.30671 (14)	0.2531 (3)	0.0175 (5)
H6	1.0859	0.3230	0.2705	0.021*
C7	0.8537 (3)	0.34266 (13)	0.3137 (3)	0.0143 (5)
H7	0.8934	0.3833	0.3711	0.017*
C8	0.3703 (3)	0.58033 (13)	0.2194 (3)	0.0143 (5)
H8	0.2668	0.5848	0.2464	0.017*
C9	0.3762 (3)	0.61133 (14)	0.0923 (3)	0.0167 (5)
H9	0.2792	0.6367	0.0354	0.020*
C10	0.5299 (3)	0.60410 (14)	0.0504 (3)	0.0159 (5)
H10	0.5376	0.6246	-0.0348	0.019*
C11	0.6709 (3)	0.56580 (13)	0.1378 (3)	0.0131 (5)
C12	0.6545 (3)	0.53690 (13)	0.2651 (3)	0.0134 (5)
H12	0.7500	0.5117	0.3243	0.016*
C13	0.8401 (3)	0.55041 (14)	0.1008 (3)	0.0152 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.01372 (14)	0.01967 (16)	0.03198 (17)	-0.00559 (9)	0.00611 (11)	-0.00455 (10)
Ni1	0.00719 (19)	0.0141 (2)	0.0102 (2)	0.00053 (14)	0.00254 (15)	0.00033 (15)
O1	0.0115 (8)	0.0168 (9)	0.0122 (8)	0.0021 (6)	0.0022 (6)	-0.0009 (6)
O2	0.0082 (7)	0.0226 (10)	0.0164 (8)	0.0018 (6)	0.0015 (6)	-0.0023 (7)
O3	0.0133 (8)	0.0269 (10)	0.0156 (9)	0.0064 (7)	0.0049 (7)	0.0048 (7)
O4	0.0083 (8)	0.0191 (10)	0.0166 (9)	0.0012 (7)	0.0024 (7)	-0.0005 (7)
N1	0.0108 (9)	0.0138 (10)	0.0142 (10)	0.0004 (7)	0.0036 (8)	-0.0001 (8)
N2	0.0131 (10)	0.0324 (13)	0.0143 (10)	0.0074 (9)	0.0063 (9)	0.0062 (9)
C1	0.0105 (10)	0.0152 (12)	0.0126 (11)	0.0013 (8)	0.0043 (9)	0.0026 (9)
C2	0.0103 (10)	0.0143 (12)	0.0143 (11)	0.0018 (8)	0.0034 (9)	0.0015 (9)
C3	0.0102 (10)	0.0162 (12)	0.0137 (11)	-0.0005 (8)	0.0013 (9)	0.0018 (9)
C4	0.0166 (11)	0.0127 (12)	0.0172 (12)	0.0027 (9)	0.0025 (9)	-0.0012 (9)
C5	0.0165 (12)	0.0207 (14)	0.0166 (12)	0.0065 (9)	0.0064 (10)	0.0025 (10)
C6	0.0113 (11)	0.0205 (13)	0.0209 (12)	-0.0004 (9)	0.0051 (10)	0.0010 (10)
C7	0.0110 (10)	0.0149 (12)	0.0165 (11)	0.0010 (8)	0.0031 (9)	0.0005 (9)
C8	0.0095 (10)	0.0163 (12)	0.0167 (11)	0.0020 (8)	0.0029 (9)	0.0001 (9)
C9	0.0093 (10)	0.0211 (13)	0.0175 (12)	0.0051 (9)	0.0002 (9)	0.0043 (10)
C10	0.0132 (11)	0.0224 (13)	0.0130 (11)	0.0022 (9)	0.0052 (9)	0.0032 (9)
C11	0.0117 (11)	0.0146 (12)	0.0139 (11)	0.0007 (8)	0.0051 (9)	-0.0004 (9)
C12	0.0117 (10)	0.0135 (12)	0.0155 (11)	0.0011 (8)	0.0046 (9)	0.0000 (9)
C13	0.0113 (11)	0.0203 (13)	0.0145 (11)	0.0005 (9)	0.0044 (9)	-0.0019 (9)

Geometric parameters (\AA , $^\circ$)

Br1—C3	1.896 (2)	C3—C4	1.390 (3)
--------	-----------	-------	-----------

Ni1—O1 ⁱ	2.0806 (16)	C4—C5	1.383 (4)
Ni1—O1	2.0806 (16)	C4—H4	0.9300
Ni1—O4	2.1012 (17)	C5—C6	1.382 (4)
Ni1—O4 ⁱ	2.1012 (17)	C5—H5	0.9300
Ni1—N1	2.068 (2)	C6—C7	1.390 (3)
Ni1—N1 ⁱ	2.068 (2)	C6—H6	0.9300
O1—C1	1.260 (3)	C7—H7	0.9300
O2—C1	1.258 (3)	C8—H8	0.9300
O3—C13	1.241 (3)	C9—C8	1.377 (3)
O4—H41	0.86 (4)	C9—H9	0.9300
O4—H42	0.83 (4)	C10—C9	1.391 (3)
N1—C8	1.347 (3)	C10—H10	0.9300
N1—C12	1.337 (3)	C11—C10	1.383 (3)
N2—H21	0.82 (4)	C11—C12	1.390 (3)
N2—H22	0.79 (4)	C12—H12	0.9300
C1—C2	1.513 (3)	C13—N2	1.327 (3)
C2—C7	1.394 (3)	C13—C11	1.505 (3)
C3—C2	1.400 (4)		
O1 ⁱ —Ni1—O1	180.000 (1)	C4—C3—Br1	115.24 (18)
O1—Ni1—O4	87.40 (7)	C4—C3—C2	121.6 (2)
O1 ⁱ —Ni1—O4	92.60 (7)	C3—C4—H4	120.2
O1 ⁱ —Ni1—O4 ⁱ	87.40 (7)	C5—C4—C3	119.7 (2)
O1—Ni1—O4 ⁱ	92.60 (7)	C5—C4—H4	120.2
O4—Ni1—O4 ⁱ	180.00 (10)	C4—C5—H5	119.9
N1—Ni1—O1	89.60 (7)	C6—C5—C4	120.2 (2)
N1 ⁱ —Ni1—O1	90.40 (7)	C6—C5—H5	119.9
N1—Ni1—O1 ⁱ	90.40 (7)	C5—C6—C7	119.5 (2)
N1 ⁱ —Ni1—O1 ⁱ	89.60 (7)	C5—C6—H6	120.3
N1—Ni1—O4	87.68 (7)	C7—C6—H6	120.3
N1 ⁱ —Ni1—O4	92.32 (7)	C2—C7—H7	119.0
N1—Ni1—O4 ⁱ	92.32 (7)	C6—C7—C2	122.0 (2)
N1 ⁱ —Ni1—O4 ⁱ	87.68 (7)	C6—C7—H7	119.0
N1—Ni1—N1 ⁱ	180.00 (10)	N1—C8—C9	123.0 (2)
C1—O1—Ni1	122.99 (15)	N1—C8—H8	118.5
Ni1—O4—H41	95 (3)	C9—C8—H8	118.5
Ni1—O4—H42	124 (2)	C8—C9—C10	118.8 (2)
H42—O4—H41	105 (4)	C8—C9—H9	120.6
C8—N1—Ni1	122.66 (15)	C10—C9—H9	120.6
C12—N1—Ni1	119.53 (16)	C9—C10—H10	120.6
C12—N1—C8	117.8 (2)	C11—C10—C9	118.8 (2)
C13—N2—H21	121 (3)	C11—C10—H10	120.6
C13—N2—H22	118 (2)	C10—C11—C12	118.7 (2)
H21—N2—H22	118 (3)	C10—C11—C13	124.0 (2)
O1—C1—C2	117.8 (2)	C12—C11—C13	117.3 (2)
O2—C1—O1	124.7 (2)	N1—C12—C11	123.0 (2)
O2—C1—C2	117.4 (2)	N1—C12—H12	118.5

supplementary materials

C3—C2—C1	124.1 (2)	C11—C12—H12	118.5
C7—C2—C1	118.8 (2)	O3—C13—N2	122.8 (2)
C7—C2—C3	117.0 (2)	O3—C13—C11	119.9 (2)
C2—C3—Br1	123.11 (17)	N2—C13—C11	117.2 (2)
O4—Ni1—O1—C1	-145.89 (18)	C1—C2—C7—C6	-172.5 (2)
O4 ⁱ —Ni1—O1—C1	34.11 (18)	C3—C2—C7—C6	2.3 (4)
N1—Ni1—O1—C1	-58.20 (18)	Br1—C3—C2—C1	-10.9 (3)
N1 ⁱ —Ni1—O1—C1	121.80 (18)	Br1—C3—C2—C7	174.53 (17)
O1—Ni1—N1—C8	137.00 (19)	C4—C3—C2—C1	171.3 (2)
O1 ⁱ —Ni1—N1—C8	-43.00 (19)	C4—C3—C2—C7	-3.2 (4)
O1—Ni1—N1—C12	-44.08 (18)	Br1—C3—C4—C5	-176.67 (19)
O1 ⁱ —Ni1—N1—C12	135.92 (18)	C2—C3—C4—C5	1.2 (4)
O4—Ni1—N1—C12	43.33 (18)	C3—C4—C5—C6	1.7 (4)
O4 ⁱ —Ni1—N1—C12	-136.67 (18)	C4—C5—C6—C7	-2.6 (4)
O4—Ni1—N1—C8	-135.59 (19)	C5—C6—C7—C2	0.5 (4)
O4 ⁱ —Ni1—N1—C8	44.41 (19)	C10—C9—C8—N1	0.7 (4)
Ni1—O1—C1—O2	-19.9 (3)	C11—C10—C9—C8	0.2 (4)
Ni1—O1—C1—C2	155.74 (16)	C12—C11—C10—C9	-0.9 (4)
Ni1—N1—C8—C9	178.11 (19)	C13—C11—C10—C9	176.4 (2)
C12—N1—C8—C9	-0.8 (4)	C10—C11—C12—N1	0.8 (4)
Ni1—N1—C12—C11	-178.89 (18)	C13—C11—C12—N1	-176.7 (2)
C8—N1—C12—C11	0.1 (4)	O3—C13—C11—C10	-173.8 (2)
O1—C1—C2—C3	156.0 (2)	O3—C13—C11—C12	3.5 (4)
O1—C1—C2—C7	-29.6 (3)	N2—C13—C11—C10	4.7 (4)
O2—C1—C2—C3	-28.0 (4)	N2—C13—C11—C12	-178.0 (2)
O2—C1—C2—C7	146.4 (2)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H21 \cdots O2 ⁱⁱ	0.82 (4)	2.09 (4)	2.864 (3)	156 (4)
N2—H22 \cdots O3 ⁱⁱⁱ	0.79 (4)	2.22 (4)	2.951 (3)	153 (4)
O4—H41 \cdots O2 ⁱ	0.86 (4)	1.77 (4)	2.612 (2)	165 (5)
O4—H42 \cdots O3 ^{iv}	0.83 (4)	2.09 (4)	2.886 (2)	162 (3)
C9—H9 \cdots Cg1 ^v	0.93	2.89	3.617 (3)	136

Symmetry codes: (ii) $-x+1, -y+1, -z$; (iii) $-x+2, -y+1, -z$; (i) $-x+1, -y+1, -z+1$; (iv) $-x+2, -y+1, -z+1$; (v) $-x, -y, -z+1$.

Fig. 1

