

Constancy of some maps into f -manifolds

Sadettin Erdem

Abstract

Constancy of some maps into certain f -manifolds is discussed

Introduction

In dealing with maps, such as harmonic and holomorphic ones, between manifolds existence questions form an essential part of their study. Mathematicians normally study such maps under certain conditions imposed on the manifolds and on the maps themselves. A vital question is that whether there exist such maps under the restrictions imposed. We highlight a few results in this line:

i) [9], If a harmonic map is constant on the boundary of a flat disk M of any dimension then it is constant on the whole disk M ;

ii) [5], Let $\phi : M \rightarrow N$ be a holomorphic map between Kaehler manifolds with M compact and $\text{rank}(\phi) < \dim(M)$. If, for the respective Kaehler forms ω^M and ω^N , the cohomology classes satisfy that $[\phi^*\omega^N] = c[\omega^M]$ for some $c \in \mathbb{R}$ then ϕ is constant. In particular, one may take M to be a complex Grassmannian or a complex quadric.

Almost a decade ago mathematicians (see for example [4], [6], [7] and [8]) started to consider holomorphic maps and harmonic maps between metric f -manifolds. The main difference between an f -manifold and a complex one is that an almost complex structure is replaced with an f -structure by which we mean a 1-1 tensor field f satisfying $f^3 + f = 0$. In 2001, the following results were established:

iii) [4], Every holomorphic map from an almost Hermitian manifold into an almost \mathcal{S} -manifold is constant.

Received by the editors October 2003.

Communicated by L. Vanhecke.

2000 *Mathematics Subject Classification* : Primary 58E20, Secondary 53C15, 53C25.

Key words and phrases : Holomorphic maps, f -manifolds.

iv) [8], Every holomorphic map from a semi-Kaehler manifold into a strongly pseudoconvex CR -manifold is constant.

In this work we give a non-existence result, *Theorem (2.1)*, for maps into certain f -manifolds under more general conditions. Our result improves the last two results above. One of the main features of our work is that it removes all restrictions on the domain manifold.

1 Preliminaries

For a Riemannian manifold $(N^{2n+\ell}, h)$ let φ denote a $(1, 1)$ - tensor field on N of rank $2n$ and nullity $\ell \geq 0$. Put $D = \varphi(TN)$ and $\mathcal{V} = \text{Ker}(\varphi)$. The distributions D and \mathcal{V} over N are called φ -horizontal and φ -vertical respectively. The triple (N, h, φ) is then called a *metric f -manifold* (M - f -manifold) provided:

- i) $\varphi^3 + \varphi = 0$;
- ii) $h(X, Y) = 0, \forall X \in D, Y \in \mathcal{V}$;
- iii) $h(X, Y) = h(\varphi X, \varphi Y) \forall X, Y \in D$.

We refer the conditions (ii) and (iii) as *h -compatibility of φ* . With no metric considered, the pair (N, φ) is called an *f -manifold*.

Suppose there is a global frame field $\{\xi_j\}_{j=1}^\ell$ for the φ -vertical bundle \mathcal{V} with dual 1-forms $\{\eta^j\}_{j=1}^\ell$ satisfying

$$\varphi^2 = (-I + \sum_{j=1}^{\ell} \eta^j \otimes \xi_j), \quad \eta^j(\xi_a) = \delta_a^j$$

and

$$h(\varphi X, \varphi Y) = h(X, Y) - \sum_{j=1}^{\ell} \eta^j(X)\eta^j(Y).$$

Then $N = (N^{2n+\ell}; h, \varphi, \xi_j, \eta^j)$ is called a *globally framed metric f -manifold*. With no metric considered, $N = (N^{2n+\ell}; \varphi, \xi_j, \eta^j)$ will be called a *globally framed f -manifold*.

For the class of globally framed metric f -manifold $N = (N^{2n+\ell}; h, \varphi, \xi_j, \eta^j)$ we list some of its subclasses for later use ([1], [2], [3], [4]):

- i) For $\ell = 0$, $N = (N^{2n}, h, \varphi)$ is called an *almost Hermitian manifold*.
- ii) For $\ell \geq 1$ and set $\Omega(X, Y) = h(X, \varphi Y)$ then
 - a $^\circ$) N is called an *almost \mathcal{S} -manifold* provided $d\eta^j = \Omega$, for each $j \in \{1, 2, \dots, \ell\}$.
 - b $^\circ$) N is called an *almost \mathcal{C} -manifold* provided $d\Omega = 0$ and $d\eta^j = 0$ for each $j = 1, 2, \dots, \ell$.

iii) If $\ell = 1$, $N = (N^{2n+1}; h, \varphi, \xi, \eta)$ is called an *almost contact metric manifold* (almost CM -manifold). If further an almost CM -manifold is also an almost \mathcal{S} -manifold (so that $d\eta = \Omega$) then we drop the adjective "almost" and simply call a *contact metric manifold* (CM -manifold). If an almost CM -manifold is also an almost \mathcal{C} -manifold, that is $d\Omega = 0$ and $d\eta = 0$, then it is called *almost cosymplectic*.

- iv) An almost CM -manifold $(N^{2n+1}, h, \varphi, \xi, \eta)$ is called
 - a $^\circ$) *nearly Sasakian* if $\forall X, Y \in \Gamma(TN)$

$$(\nabla_X \varphi)Y + (\nabla_Y \varphi)X = -2h(X, Y)\xi + \eta(X)Y + \eta(Y)X$$

where ∇ denotes the Levi-Civita connection;

b°) almost trans-Sasakian of type (α, β) if

$$d\eta = \alpha\Omega - \frac{1}{n}\eta \wedge \varphi^*(\delta\Omega) \quad \text{and} \quad d\Omega = \Omega \wedge \left(\frac{1}{n}\varphi^*(\delta\Omega) - \beta\eta\right)$$

for some functions α, β on N , where δ is the codifferential operator and $\varphi^*(\delta\Omega)(X) = (\delta\Omega)(\varphi X)$. In particular, if $\alpha = \frac{1}{2n}(\delta\Omega)(\xi)$ and $\beta = \frac{1}{n}\delta\eta = \text{div}(\xi)$ then N is simply called an almost trans-Sasakian manifold;

c°) nearly cosymplectic if $(\nabla_X \varphi)X = 0$ for all $X \in \Gamma(TN)$;

d°) quasi-K- cosymplectic if

$$\mathcal{S}(X, Y) := (\nabla_X \varphi)Y + (\nabla_{(\varphi X)} \varphi)(\varphi Y) = \eta(Y)\nabla_{(\varphi X)} \xi \quad \text{for all } X, Y \in \Gamma(TN);$$

e°) cosymplectic if it is almost cosymplectic and normal (i.e. $N^1 = [\varphi, \varphi](X, Y) + 2d\eta(X, Y)\xi = 0$ for all $X, Y \in \Gamma(TN)$).

Remark:

i) [10], Every almost cosymplectic manifold is quasi K-cosymplectic.

ii) [1], Every almost contact metric manifold $(N^{2n+1}, h, \varphi, \xi, \eta)$ is cosymplectic if and only if $\nabla\varphi = 0$, that is φ is parallel.

Let $\phi : (M, g) \rightarrow (N, h, \varphi)$ be a smooth map of a Riemannian manifold into a M - f -manifold . Set $\mathcal{K} = \mathcal{K}_\phi = \ker(d\phi)$, $\mathcal{H} = \mathcal{H}_\phi = \mathcal{K}^\perp$ and $\overline{M} = \{p \in M : d\phi_p \neq 0\}$. If \mathcal{K} (and therefore \mathcal{H}) forms a bundle then \mathcal{K} and \mathcal{H} are called the *vertical* and *horizontal distributions associated with ϕ* respectively.

Throughout our work the map ϕ will be smooth and \overline{M} will be a dense open subset of M

Definition (1.1): A smooth map $\phi : M \rightarrow (N, \varphi)$ into an f -manifold is said to be

i) φ -invariant if $d\phi(TM)$ is invariant under φ , that is, $\varphi \circ d\phi(TM) = d\phi(TM)$.

ii) properly φ -invariant if it is φ -invariant of constant rank.

Remark: Note here that

a°) the φ -invariance of ϕ implies that $d\phi(TM) \subseteq D_\varphi$. Also every φ -invariant map into a surface is necessarily a submersion on \overline{M} ;

b°) ϕ is properly φ -invariant if and only if the pull-back $\phi^{-1}(d\phi(TM)) \rightarrow \overline{M}$ together with $\phi^{-1}\varphi$ forms a complex bundle (and therefore $\mathcal{K} \rightarrow \overline{M}$ forms a bundle) over \overline{M} .

A smooth map $\phi : (M, \varphi^M) \rightarrow (N, \varphi^N)$ between f -manifolds is said to be (φ^M, φ^N) -holomorphic [resp: (φ^M, φ^N) -antiholomorphic] if

$$d\phi \circ \varphi^M = \varphi^N \circ d\phi, \quad [\text{resp: } d\phi \circ \varphi^M = -\varphi^N \circ d\phi].$$

We write $\pm(\varphi^M, \varphi^N)$ -holomorphic to mean either (φ^M, φ^N) -holomorphic or (φ^M, φ^N) -antiholomorphic.

For a φ -invariant map $\phi : (M, g) \rightarrow (N, \varphi)$ of a Riemannian manifold into an f -manifold (not necessarily of constant rank), set $\forall p \in \overline{M}$,

$$\Psi_p(X_p) = \Psi_p^\phi(X_p) = \begin{cases} (d\phi)^{-1} \circ \varphi \circ d\phi(X_p) & , \quad X_p \in \mathcal{H}_p \\ 0 & , \quad X_p \in \mathcal{K}_p \end{cases} \quad (1.1)$$

where $d\phi^{-1} = (d\phi|_{\mathcal{H}})^{-1}$.

Observe that for every φ -invariant map $\phi : (M, g) \rightarrow (N, \varphi)$ we have $d\phi \circ \Psi^\phi = \varphi \circ d\phi$. If ϕ is properly φ -invariant then Ψ becomes of constant rank and therefore it becomes an f -structure on M which we call ϕ -associated f -structure. In that case, we have that $\mathcal{H}_\phi = \mathcal{D}_\Psi$ and therefore we shall be using the letters \mathcal{H} and \mathcal{D} interchangeably for the same bundle. Setting $2m = \text{rank}(\mathcal{H})$ and $s = \text{rank}(\mathcal{K})$ we see that (M^{2m+s}, Ψ) becomes an f -manifold. Thus, every properly φ -invariant map is (Ψ, φ) -holomorphic as a map $\phi : (M, \Psi) \rightarrow (N, \varphi)$ between f -manifolds. However, this ϕ -associated f -structure Ψ need not be compatible with the prescribed metric g and therefore the triple (M^{2m+s}, g, Ψ) need not to be a M - f -manifold.

Definition (1.2): A map $\phi : (M, g) \rightarrow (N, h, \varphi)$ is said to be

a°) *horizontally weakly conformal* if $d\phi$ is surjective and satisfies that

$$h(d\phi(X), d\phi(Y)) = \lambda g(X, Y); \quad \forall X, Y \in \mathcal{H}$$

for some smooth function $\lambda : M \rightarrow \mathbb{R}$

b°) (g, φ) -pseudo horizontally weakly conformal (or simply pseudo horizontally weakly conformal when no confusion arises) if ϕ is φ -invariant and Ψ^ϕ is compatible with the metric g .

Remark: Note here that

i) every horizontally weakly conformal map is also pseudo horizontally weakly conformal,

ii) the pseudo horizontal weak conformality of ϕ does not depend on the metric h .

2 Constancy of certain maps

For a smooth map $\phi : M \rightarrow (N, \varphi)$ of a manifold into an f -manifold with $d\phi(TM) \subseteq \mathcal{D}_\varphi$ we introduce the following two conditions: For every $p \in \overline{M}$, writing $q = \phi(p)$.

\mathbf{C}_N : there is a nonzero vector $\zeta_q \in \mathcal{V}_\varphi(q) = \ker(\varphi_q)$ and a local 1-form ω defined on an open subset \mathcal{U} of N containing q with $\omega(\zeta_q) = 1$ such that $\forall X \in T_q N$

$$\varphi^2(X) = -X + \omega(X)\zeta_q + v_x \quad \text{for some } v_x \in \mathcal{V}(q) \cap \ker(\omega_q);$$

\mathbf{C}_ϕ : for $d\phi(T_p M) \neq 0$, the function $\gamma_q : d\phi(T_p M) \subset T_q N \rightarrow \mathbb{R}$ given by, (on an open subset $\mathcal{U} \subset N$ containing q) by

$$\gamma_q(X_q) = d\omega(X_q, \varphi X_q)$$

is not zero.

Observe that, one can easily deduce from the condition \mathbf{C}_N that $\mathcal{D}_\varphi(q) \subseteq \ker(\omega_q)$.

Note that, for a smooth map $\phi : M \rightarrow (N, \varphi)$, one has wide variety of choices of structures on N for which the conditions \mathbf{C}_N and \mathbf{C}_ϕ are satisfied. We give the

following examples. Let $\phi : M \rightarrow N$ be a smooth map into a M - $f.pk$ - manifold. Consider the following cases:

1°) Let $N = (N^{2n+\ell}, h, \varphi, \xi_j, \eta^j)$ be an almost \mathcal{S} -manifold or, in particular, a CM -manifold then \mathbf{C}_N and \mathbf{C}_ϕ are satisfied. Indeed, putting $\zeta = \xi_j$ and $\omega = \eta^j$ for some $j \in \{1, \dots, \ell\}$, $\ell \geq 1$ we see that

$$\mathbf{C}_N : \varphi^2(X) = -X + \omega(X)\xi_j + v \quad \text{with} \quad v = \sum_{\substack{t=1 \\ t \neq j}}^{\ell} \eta^t(X)\xi_t \in \mathcal{V} \cap \ker(\omega)$$

$\mathbf{C}_\phi : \gamma(X) = d\eta^j(X, \varphi X) = \Omega(X, \varphi X) = -h(X, X) \neq 0, \quad \forall X \in d\phi(TM) \subseteq D_\varphi$, where $\Omega(X, Y) = d\eta^j(X, Y) = h(X, \varphi Y)$.

2°) Let $N = (N^{2n+1}, h, \varphi, \xi, \eta)$ be a nearly Sasakian manifold then \mathbf{C}_N and \mathbf{C}_ϕ are satisfied. For let $\zeta = \xi$ and $\omega = \eta$, then \mathbf{C}_N is obvious. For \mathbf{C}_ϕ , since N is nearly Sasakian, we have

$$(\nabla_x \varphi)X = -h(X, X)\xi + \eta(X)X$$

But then, for $X \in \Gamma(d\phi(T\bar{M})) \subseteq \Gamma(D^N)$, this gives $(\nabla_x \varphi)X = -h(X, X)\xi$ and therefore we get

$$\mathcal{S}(X, X) = -2h(X, X)\xi \tag{2.1}$$

where $\Gamma(V)$ denotes the set of all smooth local sections of a bundle $V \rightarrow N$. On the other hand, for $X \in \Gamma(d\phi(TM))$ we have

$$\begin{aligned} d\eta(X, \varphi X) &= -\eta([X, \varphi X]) = -\eta(\nabla_x(\varphi X) + \nabla_{(\varphi X)}(\varphi^2 X)) \\ &= -\eta((\nabla_x \varphi)X + (\nabla_{(\varphi X)} \varphi)(\varphi X)) = -\eta(\mathcal{S}(X, X)) \end{aligned}$$

So, by (2.1) we get

$$\gamma(X) = 2h(X, X) \neq 0.$$

3°) Let $N = (N^{2n+1}, h, \varphi, \xi, \eta)$ be an almost trans-Sasakian manifold of type (α, β) with $\alpha(q) \neq 0, \quad \forall q \in \phi(M) \subseteq N$. Then \mathbf{C}_N and \mathbf{C}_ϕ are satisfied. For let $\zeta = \xi$ and $\omega = \eta$, then \mathbf{C}_N is obviously satisfied. For \mathbf{C}_ϕ , since N is trans-Sasakian, we have

$$d\eta(X, \varphi X) = \alpha\Omega(X, \varphi X) - \frac{1}{n}(\eta \wedge \varphi^*(\delta\Omega))(X, \varphi X) = \alpha\Omega(X, \varphi X) = -\alpha h(X, X)$$

for all $X \in \Gamma(d\phi(T\bar{M})) \subseteq \Gamma(D^N)$. Thus $\gamma(X) = -\alpha h(X, X) \neq 0$ on $d\phi(T\bar{M})$.

However, for the following structure on N , the condition \mathbf{C}_ϕ , for example, fails to hold:

Let $\phi : M \rightarrow (N^{2n+\ell}, h, \varphi, \xi_j, \eta^j)$ be a smooth map into a M - $f.pk$ -manifold N such that $\mathcal{S}(X, X) \in D^N$ and $d\phi(TM) \subseteq D^N, \quad \forall X \in D^N$ (or equivalently, $[X, \varphi X] \in D^N, \quad \forall X \in D^N$). Then $\gamma(X) = 0$ on D^N and therefore on $d\phi(TM)$. Indeed, setting $\omega = \eta^j$, for any $j = 1, 2, \dots, \ell$ note that

$$\begin{aligned} \gamma(X) &= d\omega(X, \varphi X) = -\omega([X, \varphi X]) = -\omega(\nabla_x(\varphi X) + \nabla_{(\varphi X)}(\varphi^2 X)) \\ &= -\omega((\nabla_x \varphi)X + (\nabla_{(\varphi X)} \varphi)(\varphi X) + \varphi(\nabla_x X + \nabla_{(\varphi X)}(\varphi X))) \\ &= \omega((\nabla_x \varphi)X + (\nabla_{(\varphi X)} \varphi)(\varphi X)) = \omega(\mathcal{S}(X, X)) = 0. \end{aligned}$$

Remark: Observe that for almost \mathcal{C} -manifolds, nearly cosymplectic and quasi K-cosymplectic manifolds the condition that $\mathcal{S}(X, X) \in D$, $\forall X \in D$ trivially holds.

Theorem (2.1): *Let $\phi : M \rightarrow (N^{2n+\ell}, \varphi)$ be a φ -invariant map from an arbitrary smooth manifold into an f -manifold, with $\ell = \text{rank}(\mathcal{V}_\varphi) \neq 0$ such that the conditions C_N and C_ϕ hold. Then ϕ is constant, that is $\overline{M} = \{p \in M : d\phi_p \neq 0\}$ is empty.*

Proof: Suppose \overline{M} is not empty, then $d\phi_{p_0} \neq 0$ for some $p_0 \in M$. By condition C_ϕ , there is $Z = d\phi_{p_0}(W)$ with $\gamma_{q_0}(Z) \neq 0$ for some $W \in T_{p_0}M$. Set $H = \mathcal{K}^\perp$ with respect to any chosen Riemannian metric on M and let $X \in \Gamma(H)$ be a local section with $d\phi(X_{p_0}) = Z$. Recalling the endomorphism $\Psi_p : T_pM \rightarrow T_pM$ defined by (1.1), (note that then we have $\varphi \circ d\phi = d\phi \circ \Psi$), observe that

$$A := [d\phi(X), \varphi d\phi(X)] = d\phi([X, \Psi X])$$

which shows that $A_{q_0} \in d\phi(T_{p_0}M) \subseteq D_\varphi(q_0)$ and so $\omega(A_{q_0}) = 0$. On the other hand,

$$0 = \omega(A_{q_0}) = \omega([d\phi(X), \varphi d\phi(X)]_{q_0}) = -d\omega(d\phi(X_{p_0}), \varphi d\phi(X_{p_0})) = -\gamma_{q_0}(Z).$$

This contradicts the choice of $Z \in d\phi(T_{p_0}M)$ with $\gamma_{q_0}(Z) \neq 0$, so the result follows.

Remark: In the above theorem it is essential that $\text{rank}(\mathcal{V}_\varphi) = \ell > 0$ as the conditions C_N and C_ϕ cannot possibly hold when $\ell = 0$. Also note that we do not impose any condition on M .

Corollary (2.2): *Let $\phi : (M, \varphi^M) \rightarrow (N, \varphi^N)$ be a $\pm(\varphi^M, \varphi^N)$ -holomorphic map between f -manifolds with $d\phi(TM) \subseteq D^N$ and suppose that the conditions C_N and C_ϕ hold. Then ϕ is constant.*

Proof: Observe that $\pm(\varphi^M, \varphi^N)$ -holomorphicity, in this case, implies that ϕ is φ -invariant. So by Theorem (2.1), ϕ is constant.

Corollary (2.3): *Let $(M, J) \rightarrow (N, \varphi)$ be a $\pm(J, \varphi)$ -holomorphic map from an almost complex manifold into an f -manifold such that the conditions C_N and C_ϕ hold. Then ϕ is constant.*

Proof: Observe that since J is an almost complex structure, $\pm(J, \varphi)$ -holomorphicity gives that $d\phi(TM) \subseteq D^N$. Then the result follows from Corollary (2.2).

We say that a M - f .pk manifold N is in \star -category if N is either nearly Sasakian or trans-Sasakian of type (α, β) with $\alpha(q) \neq 0$, $\forall q \in N$ or an almost \mathcal{S} -manifold.

Corollary (2.4): *Let $\phi : (M, J) \rightarrow (N^{2n+\ell}, h, \varphi, \xi_j, \eta^j)$ be a $\pm(J, \varphi)$ -holomorphic map from an almost complex manifold into a manifold which is in \star -category. Then ϕ is constant.*

Proof: Since the target manifold is in \star -category, the conditions C_N and C_ϕ are satisfied. Thus the result follows from Corollary (2.3).

In particular, when $(N^{2n+\ell}, h, \varphi, \xi_j, \eta^j)$ is an almost \mathcal{S} -manifold, Corollary (2.4) gives immediately the result in ([4], Theorem (5.2) and therefore Theorem (5.1)).

Corollary (2.5): Let $\phi : (M, g) \rightarrow (N^{2n+\ell}; h, \varphi, \xi_j, \eta^j)$ be a φ -pseudo horizontally weakly conformal map from an arbitrary Riemannian manifold into a manifold which is in \star -category, then ϕ is constant.

Proof: The φ -pseudo horizontal weak conformality of ϕ and N being in the \star -category imply that the hypothesis of Theorem(2.1) are satisfied. Thus the result follows immediately.

Following the terminology in [8], a strongly pseudoconvex CR -manifold $N = (N^{2n+1}; \varphi_D, D^N, \eta)$ with its Levi distribution D^N of rank $2n$, 1-form η and positive definite Levi form $L(X, Y) = -d\eta(\varphi_D X, Y)$ may be viewed as a contact metric manifold $(N^{2n+1}; h, \varphi, \xi, \eta)$ with $\varphi|_{D^N} = \varphi_D$, $h|_{D^N} = L$ and η its contact form, ξ its characteristic vector field. Thus for such manifolds we have :

Corollary (2.6): Any $\pm(J, \varphi)$ -holomorphic map $\phi : (M, J) \rightarrow N$ from an almost complex manifold (or any φ -pseudo horizontally weakly conformal map $\phi : (M, g) \rightarrow N$ from an arbitrary Riemannian manifold) into a strongly pseudoconvex almost CR -manifold is constant.

This Corollary recovers the result given in ([8], Proposition (2.5)). Note here that, in our work, the condition imposed therein that M is semi-Kaehler is removed. We also include the cases where ϕ is (J, φ) - antiholomorphic.

Acknowledgement: I would like to thank to the referee for all her/his suggestions and corrections.

References

- [1] Blair, D.: Riemannian Geometry of Contact and Symmetric Manifolds. *Progress in mathematics (Boston, Mass.) (2002). ISBN 0*8176-4261-7.*
- [2] Blair, D.; Showers, D. K.; Yano, K.: Nearly Sasakian structures. *Kōdai Math. Sem. Rep. 27(1976), 175-180.*
- [3] Blair, D.; Oubina, J. A.: Conformal and related changes of metric on the product of two almost contact metric manifolds. *Publ. Mat., Barc. 34 (1990), No.1, 199-207.*
- [4] Duggal, K. L.; Ianus, S.; Pastore, A. M.: Maps interchanging f -structures and their harmonicity. *Acta. Appl. Math. 67, no 1 (2001), 91-115.*
- [5] Eells, J.; Lemaire, L.: Selected topics in harmonic maps. *CBMS Regional Conference, Ser. Math., 50, Amer. Math. Soc., Providence, RI, 1983.*
- [6] Erdem, S.: φ -Pseudo harmonic morphisms, some subclasses and their liftings to a tangent bundles. *Houston J. Math, Vol.30 (2004), no. 4*
- [7] Erdem, S.: Harmonicity of maps between (indefinite) metric- f - manifolds and φ -pseudo harmonic morphisms., *Publ. Math. Debrecen, Vol.63(2003), no. 3, 317-341.*
- [8] Gherghe, C.; Ianus, S.; Pastore, A. M.: CR -manifolds, harmonic maps and stability. *J. Geom. 71(2001), 42-53 .*

- [9] Karcher, H.; Wood, J. C.: Non-existence results and growth properties for harmonic maps and forms. *J. Reine. Angew. Math.* 353 (1984), 165–180.
- [10] Olszak, Z.: On almost cosymplectic manifolds. *Kōdai Math. J. vol.4* (1981), no 2, 239-250.

Matematik Bolumu
Fen Fakultesi
Anadolu Universitesi
Eskisehir, Turkey
e-mail : s2erdem@anadolu.edu.tr